



# ARENA DISINFECTANT

## HUSKY® 891

A neutral, concentrated disinfectant cleaner that effectively cleans, deodorizes, and disinfects. Available in a wide variety of pack types and delivery systems, this product is ideal for use in educational or institutional facilities.

## Directions

1. To disinfect inanimate, hard non-porous surfaces add 1 ounce of this product per gallon of water.
2. Apply solution with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes or 1 minute for SARS-Related Coronavirus 2 (SARS-CoV-2), then remove excess liquid.
3. For sprayer applications, spray 6-8 inches from surface. Do not breathe spray mist.

**Note:** For visibly soiled areas, a pre-cleaning step is required. Surfaces which may contact food must be rinsed thoroughly with potable water after use.



**Kills SARS-CoV-2 in 60 Seconds**

## Specs

EPA Reg. No: 1839-166-8155  
 Dilution: 1:128  
 pH Level: 6.2-7.5  
 Color: Green  
 Fragrance: Fresh  
 Pack Types: Squeeze & Pour, Gallon, 5 Gallon, Cartridge, iDispense

## Features

- This product is a no rinse disinfectant cleaner that disinfects, cleans and deodorizes in one labor saving step<sup>^</sup>,2
- Kills 99.9%<sup>^</sup> of Viruses\* & Bacteria
- Fungicide (against pathogenic fungi)

<sup>^</sup>When used according to disinfection directions  
 2 Non-food contact use sites

## Recommended For

- Finished Floors
- Operating Rooms
- Chairs & Tables
- Restroom Fixtures
- Glazed porcelain, sealed marble, sealed granite
- Large Inflatable Non-Porous Plastic Structures: animals, promotional items
- Kennels & Cages



# ARENA DISINFECTANT

## DISINFECTANT CLAIMS:

### BACTERICIDAL ACTIVITY:

- Ampicillin resistant *Acinetobacter baumannii*
- Bactrim resistant *Acinetobacter baumannii*
- *Bordetella bronchiseptica*
- Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA) (NRS 123 Genotype USA400)
- Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA)
- *Corynebacterium ammoniagenes*
- *Enterobacter aerogenes*
- *Enterobacter cloacae*
- *Enterococcus faecalis*
- *Escherichia coli*
- *Fusobacterium necrophorum*
- Gentamicin resistant *Acinetobacter baumannii*
- *Klebsiella pneumoniae* subsp. *pneumoniae*
- *Lactobacillus casei* subsp. *rhamnosus*
- *Listeria monocytogenes*
- Levofloxacin resistant *Acinetobacter baumannii*
- Methicillin resistant *Staphylococcus aureus* (MRSA)
- *Pasteurella multocida*
- *Pseudomonas aeruginosa*
- *Salmonella enterica*
- *Serratia marcescens*
- *Shigella dysenteriae*
- *Shigella flexneri* Type 2b
- *Shigella sonnei*
- *Staphylococcus aureus*
- *Staphylococcus epidermidis*
- *Streptococcus pyogenes*
- Tobramycin resistant *Acinetobacter baumannii*
- Vancomycin intermediate resistant *Staphylococcus aureus* (VISA)
- Vancomycin resistant *Enterococcus faecalis* (VRE)
- *Xanthomonas maltophilia*

### VIRUCIDAL ACTIVITY:

- Avian Influenza A Virus
- Bovine Rhinotracheitis Virus
- Bovine Viral Diarrhea Virus (BVDV)
- Canine Distemper Virus
- Feline Leukemia Virus
- Feline Picornavirus
- Hepatitis B virus (HBV)
- Hepatitis C virus (HCV)
- Herpes Simplex Type 1 Virus
- Herpes Simplex Type 2 Virus
- HIV-1
- Human Coronavirus
- Influenza A2/Hong Kong Virus
- Paramyxovirus
- Porcine Respiratory & Reproductive Syndrome Virus (PRRSV)
- Rabies Virus
- Rotavirus
- SARS associated Coronavirus
- SARS-Related Coronavirus 2
- Vaccinia Virus

### FUNGICIDAL ACTIVITY:

- *Trichophyton interdigitale*, formerly known as *Trichophyton mentagrophytes* (Athletes Foot Fungus) (a cause of ringworm) (a cause of ringworm of the foot)

This is not a complete list of organisms and names