

Peroxall TB is a hydrogen peroxide based one minute acting disinfectant⁶ and is effective against SARS-CoV-2¹ virus, the cause of COVID-19. Kills Candida auris on hard, non-porous surfaces. Peroxall TB destroys 99.9% of bacteria, leaving surfaces¹ clean and disinfected and deodorizes by killing bacteria that cause offensive odors.

Peroxall TB Kills Bacteria and Viruses*. Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus, Candida auris in 10 minutes, *HIV-1, *SARS-Related Coronavirus 2, *Canine Parvovirus, *Feline Calicivirus, on environmental surfaces¹ in just 1 minute⁶
AREAS OF USE: Use Peroxall TB on surfaces¹ in public areas, hospitals, schools, institutional facilities, universities, airports, cruise ships, dental offices, veterinary clinics, rehabilitation centers, long term care facilities, athletic facilities, hotels and restaurants. Use this product on washable hard, nonporous surfaces of: Autoclaves, Cabinet Handles, Computer Peripherals, Diagnostic Equipment, Equipment Surfaces, Equipment Tables, Exam Tables, High Touch Surfaces, Infant Incubators, Instrument Stands, Lab Benches, Medical Scales, Microscope, Rescue Tools, Stethoscopes, Other hard nonporous surfaces made of: Acrylic, Formica®, Glass, Glazed Ceramic, Glazed Enameled Surfaces, Glazed Porcelain, Metal, Plastic Such as Polystyrene or Polypropylene, Stainless Steel, Upholstery-vinyl and Plastic, Power Cords, PVC Tubing, Remote Controls, Thermometers, Transport Monitors, Ultra-sound Transducers And Equipment, Walls, Animal Equipment, Animal Crates, Automatic Feeders, Buckets, Brushes, Cages, Counters, Desks, Dog Runs, Examination Tables, External Surfaces of Veterinary Equipment and Watering Appliances, Pens, Stalls.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **GENERAL DISINFECTION / *VIRUCIDAL / Candida auris DIRECTIONS:** Spray 6 - 8 inches from hard, non-porous surface, until non-porous surface surfaces are thoroughly wet. Do not breathe spray. Treated non-porous surface surfaces must remain wet for 1 minute⁶. For SARS-CoV-2, treated surfaces must remain visibly wet for 30 seconds. Wipe dry with a clean cloth, sponge, mop or towel, or allow to air dry. This product is not to be used as a terminal sterilant / high level disinfectant on any surface or instrument that is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Special Instructions for Cleaning Prior to Disinfection against Candida auris

Personal Protection: Wear appropriate barrier protection such as gloves, gowns, masks, or eye covering. **Cleaning Procedure:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the product. Pre-cleaning is to include vigorous wiping and/or scrubbing and all visible soil is removed. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading the organism. Restrooms are to be cleaned last. Do not reuse soiled cloths. **Infectious Waste Disposal:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

To Use as a One-Step¹ Disinfectant⁴: 1. Pre-clean visibly soiled areas. 2. Spray 6 – 8 inches from surface¹, making sure to wet thoroughly. 3. Allow surface to remain wet for 1 minute⁶. For SARS-CoV-2, treated surfaces¹ must remain visibly wet for 30 seconds. 4. Wipe surface dry. 5. Allow to air dry.

Manufactured By:
CANBERRA CORPORATION
3610 N. Holland-Sylvania Rd.
Toledo, OH 43615
TF (800) 832-8992

***KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES¹ / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS). Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

Special Instructions for Cleaning and Decontamination against HIV-1 on Surfaces / Objects Soiled With Blood / Body Fluids. Personal protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection. **Cleaning procedure:** Blood and other body fluids containing HIV must be thoroughly cleaned from surfaces and objects before application of Peroxall TB. Peroxall TB can be used for this purpose. **Contact time:** Leave hard, non-porous surface to remain visibly wet for 1 minute for HIV-1. Use appropriate contact time as listed on label for all other viruses, and bacteria claimed. **Disposal of infectious material:** Blood, body fluids, cleaning materials and clothing must be disposed of according to local regulations for infectious waste disposal.

Animal Housing Facilities Including Veterinary Clinics, Equine Farms, Livestock, Swine and Poultry Houses: For cleaning and disinfecting the following surfaces¹: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc. 1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. 2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. 3. Empty or cover all troughs, racks, and other feeding and watering appliances. 4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. 5. Saturate all surfaces by spraying with product until thoroughly wet. To disinfect, all surfaces¹, must remain wet for 1 minute⁶. 6. Immerse or spray all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. 7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. 8. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Open dumping is prohibited. Store in original container in areas inaccessible to children. **Container Disposal:** Nonrefillable container. Do not reuse or refill this container. **Plastic Containers:** Triple rinse or equivalent. Offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Formica® is a registered trademark of the Diller Corporation

End Notes:

- ¹ On hard, non-porous, non-food surfaces
- ⁴ When directions for disinfection are followed
- ⁶ 5 minute contact time is required for Canine Parvovirus efficacy claim, 10 minute contact time is required for Candida auris efficacy claim

110724
Label#1293



Peroxall™ TB

845

Hydrogen Peroxide
Disinfectant

Tuberculocide

Clean and disinfect in
just 1 minute⁶

EPA REG. NO. 6836-442-8155
EPA EST. NO. 8155-OH-1

NET CONTENTS:
One Quart (32 fl. oz.) 946 ml

ACTIVE INGREDIENTS:

Hydrogen Peroxide.....1.24%

OTHER INGREDIENTS:.....98.76%

TOTAL:.....100.00%

**KEEP OUT OF REACH OF
CHILDREN
CAUTION**

FIRST AID: IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

