



# Peroxall™ TB

## HUSKY® 845

Ready-to-use hydrogen peroxide disinfectant powered by proprietary HydrOx+ Technology. This one-step formula delivers broad-spectrum disinfection with a 1-minute contact time against organisms including *Mycobacterium bovis* (TB), Norovirus, MRSA, and SARS-CoV-2. Also effective against Canine Parvovirus (5 minutes) and *Candida auris* (10 minutes). Ideal for hospitals, healthcare, long-term care, schools, veterinary, and commercial settings. Peroxall TB provides hospital-grade disinfection with a Caution signal word. No dilution or mixing, just spray and wipe.

## Directions

### To Use as a One-Step<sup>1</sup> Disinfectant<sup>4</sup>:

1. Pre-clean visibly soiled areas.
2. Spray 6 – 8 inches from surface<sup>1</sup>, making sure to wet thoroughly.
3. Allow surface to remain visibly wet for 1 minute<sup>6</sup>. For SARS-CoV-2, treated surfaces must remain visibly wet for 30 seconds.
4. Wipe surface dry.



Broad-spectrum disinfection for everyday use.

## Specs

**Registrations:** EPA Registered  
**EPA Reg. No:** 6836-442-8155  
**Dilution:** RTU  
**pH Level:** 0.5–1.5  
**Color:** Clear  
**Fragrance:** Citrus  
**Pack Size:** Quart

## Features

- Cleans and disinfects in one step
- Convenient<sup>6</sup> 1 minute hospital disinfectant
- Kills 99.9% of bacteria
- Free from bleach, peracetic acid, and alcohols
- Fresh citrus scent, no lingering residue or strong fumes
- Time-saving, cost-saving formula

## Recommended For

- Acrylic
- Finished Woodwork
- Formica®
- Glass
- Glazed Ceramic & Porcelain
- Metal Plastic
- Sealed Granite,
- Marble & Quartz
- Stainless Steel
- Upholstery – Vinyl & Plastic
- Washable Hard, Non-Porous Surfaces



# Peroxall™ TB

## DISINFECTANT CLAIMS

### Bacteria

- Acinetobacter baumannii – 1 min
- Corynebacterium bovis – 1 min
- Enterobacter aerogenes
- Enterobacter cloacae New Delhi Metallo-Beta-Lactamase 1 (NDM-1) – 1 min
- Enterococcus faecalis (Vancomycin Resistant) (VRE) – 1 min
- Enterococcus faecium – 1 min
- Escherichia coli – 1 min
- Escherichia coli O157:H7 – 1 min
- Klebsiella pneumoniae – 1 min
- Klebsiella pneumoniae (Carbapenem Resistant) – 1 min
- Klebsiella pneumoniae New Delhi Metallo-Beta-Lactamase 1 (NDM-1) – 1 min
- Mycobacterium bovis BCG (TB) at 21 °C – 1 min
- Pseudomonas aeruginosa – 1 min
- Salmonella enterica – 1 min
- Staphylococcus aureus – 1 min
- Staphylococcus aureus (Methicillin Resistant) (MRSA) – 1 min
- Staphylococcus aureus (Vancomycin Resistant) (VRSA) – 1 min

### Fungi

- Candida auris – 10 min

### Viruses\*

- \*Adenovirus Type 5 – 1 min
- \*Hepatitis B Virus (HBV) – 1 min
- \*Hepatitis C Virus (HCV) – 1 min
- \*Herpes simplex type 1 Virus (HSV-1) – 1 min
- \*Human immunodeficiency virus type 1 (HIV-1) – 1 min
- \*Human Rotavirus – 1 min
- \*Influenza A Virus – 1 min
- \*Norovirus – 1 min
- \*Rhinovirus type 37 – 1 min
- \*SARS-Related Coronavirus 2 (SARS-CoV-2) (cause of COVID-19) – 30 seconds

### Animal Viruses\*

- \*Canine Parvovirus – 5 min
- \*Feline Calicivirus – 1 min
- \*Minute virus of mice – 5 min

<sup>1</sup>On hard, non-porous, non-food surfaces

<sup>4</sup>When directions for disinfection are followed

<sup>6</sup>5-minute contact time required for Canine Parvovirus efficacy claim; 10 minutes for Candida auris; 5 minutes for Minute Virus of Mice